

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Redefining the treatment paradigm for multiple myeloma

This is a pre print version of the following article:

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1732446> since 2021-02-01T12:06:44Z

Published version:

DOI:10.1016/S1470-2045(19)30295-5

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)



UNIVERSITÀ DEGLI STUDI DI TORINO

*This is an author version of the following contribution:
Questa è la versione dell'autore dell'opera:*

Gay, F., Mina, R.

16315707000;55382956600;

Redefining the treatment paradigm for multiple myeloma

(2019) The Lancet Oncology, 20 (6), pp. 743-744.

<https://www.scopus.com/inward/record.uri?eid=2-s2.0-85067196653&doi=10.1016%2fS1470-2045%2819%2930295-5&partnerID=40&md5=5a90d1fd2f96182d72e78dd892e892f6>

DOI: 10.1016/S1470-2045(19)30295-5

PUBMED ID: 31097404

DOCUMENT TYPE: Note

PUBLICATION STAGE: Final

SOURCE: Scopus

The definitive version is available at:

La versione definitiva è disponibile all'URL:

[https://www.thelancet.com/journals/lanonc/article/PIIS1470-2045\(19\)30295-5/fulltext](https://www.thelancet.com/journals/lanonc/article/PIIS1470-2045(19)30295-5/fulltext)

Continuous immunomodulation: Redefining the treatment paradigm of multiple myeloma?

Francesca Gay¹ and Roberto Mina¹

¹ Myeloma Unit, Division of Hematology, University of Torino, Azienda Ospedaliero-Universitaria Città della Salute e della Scienza di Torino, Torino, Italy.

Correspondence to: Francesca Gay, MD, Myeloma Unit, Division of Hematology, University of Torino, Azienda Ospedaliero-Universitaria Città della Salute e della Scienza di Torino, Via Genova 3, 10126 - Torino, Italy; e-mail: fgay.cittadellasalute@gmail.com; tel.: +390116334279.

In recent years, we witnessed a dramatic shift in the treatment paradigm of multiple myeloma (MM). The upfront strategy in transplant-eligible MM patients now often includes a 3-drug induction combining a proteasome inhibitor (PI; mainly bortezomib) and immunomodulatory-drug (IMiD), followed by autologous stem-cell transplant (ASCT) and lenalidomide maintenance.¹ In the non-transplant setting, continuous lenalidomide-dexamethasone (Rd) is a standard regimen, and currently the backbone of several 3-drug combinations evaluated in clinical trials (in combination with bortezomib or daratumumab).² Newly diagnosed MM patients exposed to lenalidomide at first-line will increase in the near future. Consequently, because lenalidomide is usually administered until progression, most of these patients become refractory to the drug at first relapse.

Nevertheless, Rd is the backbone of several 3-drug regimens at first relapse, combined with either a PI or a monoclonal antibody (MoAb).³ To date, no data are available on the efficacy of lenalidomide-based combinations in lenalidomide-refractory patients. Thus, with the constant increase of lenalidomide-refractory patients at first relapse, the current recommendations for treatment at relapse need to be re-defined.

Pomalidomide, a third-generation IMiD more potent than lenalidomide, has been approved in combination with dexamethasone (Pd) in patients who had failed prior lenalidomide and a PI. Based on current data in heavily pretreated patients, only one-third of patients receiving Pd ultimately reach an objective response (ORR).⁴ Many attempts to build upon Pd have been made, with several early-phase (I/II) studies focusing on lenalidomide-exposed and/or refractory patients, with promising preliminary efficacy data.

OPTIMISMM⁵ is the first randomized, phase 3 study to investigate pomalidomide early in the course of the disease, showing that a 3-drug combination including an IMiD and a PI is better than a 2-drug regimen. Of note this advantage does not come at the cost of increased toxicity; also, using pomalidomide precociously, ideally at first relapse, when the disease is more sensitive and the bone marrow microenvironment less compromised, prolongs progression-free survival (PFS; median, 21 months) compared to its use in later lines. Thus, Pvd stands out as an effective, safe and potentially cheap option (due to the expiration of bortezomib patent in the next future) at first relapse. Importantly, the combination was effective also in high-risk patients, who currently represent an unmet medical need.

Still, some open questions remain. Several compounds have been so far combined with pomalidomide in phase I/II studies, including alkylators,⁶ PIs (carfilzomib and ixazomib)^{7,8} and MoAbs (anti-CS1 and anti CD38),^{9,10} all showing promising ORR (50-86%) and prolonged PFS (median, 8.2-10.3 months). The advantage of combining pomalidomide with bortezomib rather than another drug still needs to be proved, particularly considering that bortezomib is already a backbone of frontline regimens. In fact, patients refractory to full dose bortezomib were excluded from the OPTIMISMM trial.

It is unlikely that a randomized trial will directly compare the various pomalidomide-based combinations; therefore, a bucket of compounds will emerge as available options, among which clinicians will have to choose the ideal partner for pomalidomide, weighing patient and disease characteristics, prior therapies and costs.

An exploratory analysis of the OPTIMISM trial showed an improved time-to-next-therapy of about 14 months, attributable to the immune-enhancing effects associated with pomalidomide, and supporting a continuous immunomodulatory treatment. Nevertheless, in the context of the clonal disease evolution of MM, the benefit of re-treatment with a more potent IMiD rather than switching to different drug classes still needs to be evaluated. In this regard, the current definition of lenalidomide-refractoriness does not consider the dose and the treatment-duration of lenalidomide at which refractoriness develops. In the present study, 70% of patients were refractory to lenalidomide; however, the proportion of those refractory to full-dose (25 mg) or to a lower, maintenance-dose (10-15 mg) is unknown. It is important to understand whether refractoriness to lenalidomide is dose and time-exposure dependent or not and whether the efficacy of pomalidomide varies according to lenalidomide-refractoriness definition. Also, a longterm exposure to lenalidomide may suggest sensitivity to IMiDs and thus justifies a subsequent treatment with pomalidomide; whereas an early refractoriness to lenalidomide might advocate for a drug-class shift.

PVd represents one of the future standards of care in lenalidomide-refractory patients, particularly at first relapse. Alongside PVd, other regimens with or without pomalidomide will emerge as alternatives at first relapse. Lacking reliable biomarkers predictive of a specific-drug disease sensitivity, a deeper knowledge of the meaning of refractoriness is necessary. This will allow clinicians to tailor treatment not only according to prognostic baseline parameters, but also based on dynamic predictive markers of disease sensitivity to a specific drug or class, ultimately improving treatment efficacy.

Authorship: FG and RM wrote the commentary.

Conflicts of interest disclosure: FG has received honoraria from Amgen, Bristol-Myers Squibb, Celgene, Janssen, Takeda, and served on the advisory boards for Amgen, Bristol-Myers Squibb, Celgene, Janssen, Roche, and Takeda; RM has received honoraria from Amgen and Janssen.

References:

- 1 Gay F, Engelhardt M, Terpos E, *et al.* From transplant to novel cellular therapies in multiple myeloma: European Myeloma Network guidelines and future perspectives. *Haematologica* 2018; **103**: 197–211.
- 2 Larocca A, Dold SM, Zweegman S, *et al.* Patient-centered practice in elderly myeloma patients: an overview and consensus from the European Myeloma Network (EMN). *Leukemia* 2018; **32**: 1697–712.
- 3 Larocca A, Mina R, Gay F, Brinchen S, Boccadoro M. Emerging drugs and combinations to treat multiple myeloma. *Oncotarget* 2017; **8**: 60656–72.
- 4 Siegel D, Schiller GJ, Song KW, *et al.* Pomalidomide (POM)+Low-Dose Dexamethasone (LoDEX) Safety and Efficacy in Patients (pts) with Relapsed and/or Refractory Multiple Myeloma (RRMM) Previously Treated with a Proteasome Inhibitor (PI) and in Whom Last Prior Therapy with Lenalidomide (LEN) Failed. *Blood* 2017; **130**: abstract 1812.
- 5 TO JOURNAL STAFF: Please insert OPTIMISM reference here.
- 6 Garderet L, Kuhnowski F, Berge B, *et al.* Pomalidomide, cyclophosphamide, and dexamethasone for relapsed multiple myeloma. *Blood* 2018; **132**: 2555–63.

- 7 Brinchen S, Mina R, Cafro AM, *et al.* Once-weekly carfilzomib, pomalidomide, and low-dose dexamethasone for relapsed/refractory myeloma: a phase I/II study. *Leukemia* 2018; **32**: 1803–7.
- 8 Krishnan A, Kapoor P, Palmer JM, *et al.* Phase I/II trial of the oral regimen ixazomib, pomalidomide, and dexamethasone in relapsed/refractory multiple myeloma. *Leukemia* 2018; **32**: 1567–74.
- 9 Dimopoulos MA, Dytfeld D, Grosicki S, *et al.* Elotuzumab plus Pomalidomide and Dexamethasone for Multiple Myeloma. *N Engl J Med* 2018; **379**: 1811–22.
- 10 Chari A, Suvannasankha A, Fay JW, *et al.* Daratumumab plus pomalidomide and dexamethasone in relapsed and/or refractory multiple myeloma. *Blood* 2017; **130**: 974–81.